

AD A124 384


CAPILLARY SURFACE CONTINUITY ABOVE IRREGULAR DOMAINS
(U) WISCONSIN UNIV-MADISON MATHEMATICS RESEARCH CENTER
N J KOREVAAR OCT 82 MRC-TSR-2439 DAAG29-80-C-0041

1/1


UNCLASSIFIED

F/G 12/1

NL



END
DATE
FILMED
DTIC



MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS-1963-A

ADA 124384

MRC Technical Summary Report #2439

CAPILLARY SURFACE CONTINUITY ABOVE
IRREGULAR DOMAINS

Nicholas J. Korevaar

Mathematics Research Center
University of Wisconsin-Madison
610 Walnut Street
Madison, Wisconsin 53706

October 1982

DTIC
ELECTE
FEB 15 1983
A

DTIC FILE COPY

(Received July 12, 1982)

Approved for public release
Distribution unlimited

Sponsored by

U. S. Army Research Office
P. O. Box 12211
Research Triangle Park
North Carolina 27709

National Science Foundation
Washington, D.C. 20550

83 02 014 124

UNIVERSITY OF WISCONSIN-MADISON
MATHEMATICS RESEARCH CENTER

CAPILLARY SURFACE CONTINUITY ABOVE IRREGULAR DOMAINS

Nicholas J. Korevaar

Technical Summary Report #2439
October 1982

ABSTRACT

There are simple domains Ω in \mathbb{R}^2 having re-entrant corners for which the variational solution u to the capillary problem exists, is bounded, but does not extend continuously to $\partial\Omega$. How possible is it to characterize domains for which u must extend continuously? This paper contains the following answer:

Theorem let $P_0 \in \partial\Omega$. Let u be the (variational) solution to the capillary problem in Ω with contact angle γ , $0 < \gamma < \pi/2$. If any of (i), (ii), or (iii) hold, then u extends continuously to P_0 :

- (i) Ω is convex in a neighborhood of P_0 , $\partial\Omega$ has local Lipschitz constant $L_1 < \tan \gamma$.
- (ii) $\partial\Omega$ is (locally) two C^1 curves meeting at P_0 with interior angle θ satisfying $\pi - 2\gamma < \theta < \pi$.
- (iii) $\partial\Omega$ is C^1 at P_0 .

AMS(MOS) Subject Classification: 35J20, 35J25.

Key Words: capillarity, boundary regularity, irregular, continuity, contact angle

Work Unit Number 1 - Applied Analysis

Sponsored by the U.S. Army under Contract No. DAAG29-80-C-0041. This material is based upon work supported by the National Science Foundation under Grant No. MCS-7927062, Mod. 1.

SIGNIFICANCE AND EXPLANATION

After water is poured into a vertical tube with horizontal cross section Ω the function u whose graph S_u describes the meniscus height satisfies a nonlinear elliptic partial differential equation: The mean curvature of S_u is proportional to its height above a fixed reference plane, and at the boundary the angle of contact between (the lower normal of) S_u and (the exterior normal of) $\partial\Omega \times \mathbb{R}$ is physically determined (constant if the cylinder is of uniform construction).

Nonlinear elliptic equations with co-normal derivative boundary conditions, of which the above capillary problem with prescribed contact angle is a prime example, occur frequently in physical problems but have not been as extensively studied as those with Dirichlet boundary conditions (u prescribed on $\partial\Omega$). There are many known results about when variational (weak) solutions to Dirichlet problems extend continuously to the boundary, but there is not much literature about the same question for co-normal derivative problems. That is the problem studied in this paper--for the special case of a two dimensional capillary surface making prescribed angle of contact with the bounding cylinder.

Roughly speaking, the main theorem in this paper says that u extends continuously to a point $P_0 \in \partial\Omega$ if $\partial\Omega$ is C^1 there, if $\partial\Omega$ is the vertex of two C^1 curves meeting with a convex (less than π) interior angle there, or if $\partial\Omega$ is convex with a not too large Lipschitz constant there. Since there are easy counterexamples to show that u need not extend continuously if P_0 is the vertex of a re-entrant corner (interior angle greater than π), these conditions are almost necessary as well as being sufficient. Because well known results show that u actually extends smoothly wherever $\partial\Omega$ is sufficiently smooth, the importance of these results is that they require a minimal amount of regularity for $\partial\Omega$.



Distribution For	
General	<input checked="" type="checkbox"/>
Lab	<input type="checkbox"/>
Announced	<input type="checkbox"/>
Classification	
Distribution/	
Availability Codes	
Avail and/or	Special
A	

The responsibility for the wording and views expressed in this descriptive summary lies with MRC, and not with the author of this report.

CAPILLARY SURFACE CONTINUITY ABOVE IRREGULAR DOMAINS

Nicholas J. Korevaar

§1. Introduction

For a Lipschitz domain Ω in \mathbb{R}^2 a function $u \in C^2(\Omega) \cap C^1(\bar{\Omega})$ is a classical solution to the capillary problem in a gravitational field if:

$$(1) \quad \operatorname{div} T_u = 2H(S_u) = \kappa u \text{ in } \Omega,$$

$$T_u = \frac{Du}{\sqrt{1 + |Du|^2}}, \quad H(S_u) = \text{mean curvature of the graph } S_u \text{ of } u, \quad \kappa > 0.$$

$$(2) \quad T_u \cdot n = \cos \gamma \text{ on } \partial\Omega,$$

$$0 < \gamma < \pi \text{ prescribed, } n = \text{exterior normal to } \partial\Omega.$$

Physically, S_u describes the capillary surface formed when a vertical cylinder with horizontal cross section Ω is placed in an infinite reservoir of liquid having rest height equal to zero.

Then

$$\kappa = \frac{\rho g}{\sigma} \quad \text{where } \rho = \text{density of liquid}$$

g = (downward) acceleration of gravity

σ = surface tension between liquid and air

$$\cos \gamma = \frac{\sigma_1}{\sigma} \quad \text{where } \sigma_1 = \text{surface attraction between liquid and cylinder.}$$

Geometrically, γ is the contact angle between the (downward normal to the) capillary surface S_u and the (exterior normal to the) bounding cylinder $\partial\Omega \times \mathbb{R}$.

The function u minimizes the variational problem associated with (1,2) if it minimizes the energy functional

$$(3) \quad E(f) = \int_{\Omega} (\sqrt{1 + |Df|^2} + \frac{\kappa}{2} f^2) - \int_{\partial\Omega} v f, \quad v = \frac{\sigma_1}{\sigma} = \cos \gamma$$

over the appropriate space of functions.

There are simple domains Ω in \mathbb{R}^2 having re-entrant corners for which the variational solution to the capillary problem exists, is bounded, but does not extend continuously to $\partial\Omega$ [13]. How possible is it to characterize domains for which u must extend continuously?

Theorem. Let $P_0 \in \partial\Omega$. Let u be the (variational) solution to the capillary problem in Ω with contact angle γ , $0 < \gamma < \pi/2$. If any of (i), (ii), or (iii) hold, then u extends continuously to P_0 :

(i) Ω is convex in a neighborhood of P_0 , $\partial\Omega$ has local Lipschitz constant $L_1 < \tan \gamma$;

(ii) $\partial\Omega$ is (locally) two C^1 curves meeting at P_0 with interior angle θ satisfying $\pi - 2\gamma < \theta < \pi$;

(iii) $\partial\Omega$ is C^1 at P_0 .

Since well known boundary regularity results imply that u extends smoothly wherever $\partial\Omega$ is smooth enough (see e.g. [17, 19]), the importance of this theorem is that it does not require much regularity for $\partial\Omega$. Roughly speaking, in order for u to have a jump at P_0 it needs room, such as at a re-entrant corner. If the corner is convex or there is no corner, a bounded u can have no jump and must be continuous.

Studying the same problem, Simon has shown that if P_0 is the vertex of two $C^{1,\alpha}$ curves meeting with interior angle θ , $\pi - 2\gamma < \theta < \pi$, then u actually extends to be C^1 at P_0 [16]. His method requires a strict corner $\theta < \pi$, $C^{1,\alpha}$ curves, and uses geometric measure theory to get a strong result: u is C^1 at P_0 . Our method requires less of the boundary, uses only comparison methods, but gives a weaker result: u is continuous at P_0 .

(The condition $\theta > \pi - 2\gamma$ (and the related $L_1 < \tan \gamma$) that occurs in these theorems is essentially geometric: If u was to extend to be C^1 at P_0 and if the contact angle condition was to be satisfied with both faces of $\partial\Omega \times \mathbb{R}$ there, then $\theta > \pi - 2\gamma$ is necessary for the tangent plane to exist. For $\theta < \pi - 2\gamma$ no tangent plane can exist, and in fact u approaches infinity as P_0 is approached in this case [3]).

§2. Preliminaries

First some notation: Domains Ω shall be bounded with Lipschitz boundary $\partial\Omega$ unless otherwise stated. We restrict to the case $\Omega \subset \mathbb{R}^2$. All integrals over a domain are with respect to (2-dimensional) Lebesgue measure, all integrals over boundaries of domains are with respect to (1-dimensional) Hausdorff measure.

$|\Omega|$ is the area of Ω , $|\partial\Omega|$ is the length of $\partial\Omega$. $B_\rho(P_0)$ and $S_\rho(P_0)$ are the open ball of radius ρ centered at P_0 and its boundary sphere.

We discuss capillary surfaces that solve the variational problem (3) associated with (1) and (2); u should minimize

$$\tilde{E}(f) = \int_{\Omega} (\sigma \sqrt{1 + |Df|^2} + \frac{\rho g}{2} f^2) - \int_{\partial\Omega} \sigma_1 f$$

over the appropriate space of functions. The three terms making up the energy functional are (in order) surface energy, potential energy from gravity, and wetting energy. Emmer [5] and Finn-Gerhardt [7] have studied the existence of variational solutions u to the capillary problem in Lipschitz domains Ω . When it exists the function u is real analytic in Ω and satisfies (1). Wherever $\partial\Omega$ is smooth enough u extends smoothly and satisfies the boundary condition (2) classically [17, 19].

Even when $\partial\Omega$ is only Lipschitz, however, the fact that u minimizes the energy (3) implies that (2) is satisfied in some weak (integral) sense. This weak boundary behavior is still strong enough to use a comparison method for surfaces of related

mean curvature and contact angle. The comparison method has been widely used in capillarity [1-4, 7, 13-15, 20, 22-24] and a version of it is also the primary tool used to prove our main theorem.

We state below the particular form of the comparison principle used here, after defining the type of weak boundary behavior for which it is valid.

Definition. A sequence of domains $\{\Omega_k\}$ exhausts Ω if each $\partial\Omega_k \in C^1$, $\bar{\Omega}_k \subset \Omega_{k+1}$ and $\bigcup_{k=1}^{\infty} \Omega_k = \Omega$.

Definition. Let $u \in C^2(\Omega)$, $v \in L^\infty(\partial\Omega)$. $Tu \cdot n = v$ weakly on $\partial\Omega$ if for any exhausting sequence $\{\Omega_k\}$, and for any $f \in L^\infty(\Omega) \cap W^{1,1}(\Omega)$,

$$\lim_{k \rightarrow \infty} \int_{\partial\Omega_k} f Tu \cdot n_k = \int_{\partial\Omega} v f \quad n_k = \text{exterior normal to } \partial\Omega_k.$$

Local boundary values of $Tu \cdot n$ must sometimes be considered. This can be done in a manner consistent with the above definition:

Definition. Let $U \subset \mathbb{R}^n$ be open. Let $u \in C^2(\Omega)$, $v \in L^\infty(\partial\Omega \cap U)$. $Tu \cdot n = v$ weakly on $\partial\Omega \cap U$ if for any exhausting sequence $\{\Omega_k\}$ of Ω and any $f \in L^\infty(\Omega) \cap W^{1,1}(\Omega)$ s.t. the essential support of the trace of f on $\partial\Omega$ is compactly contained in $U \cap \partial\Omega$,

$$\lim_{k \rightarrow \infty} \int_{\partial\Omega_k} f Tu \cdot n_k = \int_{\partial\Omega} v f.$$

Remark. To prove that the variational solution satisfies $Tu \cdot n = v$ weakly, one does the usual Euler-Lagrange derivation of equations and boundary conditions from (3), except that one integrates by parts over Ω_k instead of over Ω (u is smooth in Ω , hence over $\bar{\Omega}_k$). The details can be found in [7], or for a more systematic exposition, [14]. In the same references it is also shown that a smooth surface S_v that classically satisfies $Tv \cdot n = v$ (wherever n is defined) also satisfies $Tv \cdot n = v$ weakly.

Comparison Principle. (Essentially Lemma 3 of [7]) Let $v, w, \in C^2(\Omega)$. Consider a component O of the set on which $v > w$. Suppose that:

- (i) $x \in O \Rightarrow \operatorname{div} Tv(x) > \operatorname{div} Tw(x)$,
- (ii) on $\bar{O} \cap \partial\Omega$, $Tv \cdot n < Tw \cdot n$ weakly.

Then $O = \emptyset$. (So if (i) and (ii) are true for all components, $v \leq w$).

(Condition (ii) means that there is some open set $U \supset \bar{O} \cap \partial\Omega$ on which $Tv \cdot n$ and $Tw \cdot n$ are defined weakly and that their weak values satisfy $Tv \cdot n < Tw \cdot n$ almost everywhere on $\bar{O} \cap \partial\Omega$). See [7] or [14] for the proof of the comparison principle.

§3. Proof of Theorem

The techniques used to prove the main theorem are barehanded. The accompanying figures should help keep notation and ideas organized. The proof follows this outline:

We consider the domain Ω , $P_0 \in \partial\Omega$, and the variational solution $u \in C^2(\Omega) \cap H^{1,1}(\Omega)$ of (3). We take γ to be constant, $0 < \gamma < \pi$. (Essentially the same proof works if γ is allowed to be a continuous function on $\partial\Omega$). Since u minimizes (3) iff $-u$ minimizes (3) with v replaced by $-v$, it is no loss of generality to assume $0 < \gamma < \pi/2$, and we do.

Let

$$U = \limsup_{P \rightarrow P_0, P \in \Omega} u(P), \quad L = \liminf_{P \rightarrow P_0, P \in \Omega} u(P).$$

If L and U are finite, then certain geometric constraints on $\partial\Omega$ imply the existence of "ridges" and "ditches": For any $\epsilon > 0$, $\exists \rho_1 > 0$ so that for any radius ρ , $0 < \rho < \rho_1$, there is a ridge R_ρ^ϵ and a ditch D_ρ^ϵ . These are connected components in $B_\rho(P_0) \cap \Omega$ of the sets on which $u(P) > U - \epsilon$ and $u(P) < L + \epsilon$, respectively, and extend from P_0 to $S_\rho(P_0)$. (See Figure 2 and Lemma 3).

For domains satisfying conditions (i), (ii), or (iii) of the main theorem an explicit comparison function is constructed (Figure 3 and Lemma 4). If $L \neq U$, it is used to slice through either the ridge or the ditch and thus to contradict the comparison principle (Figures 3, 4, and Theorem 1). Thus, $L = U$ and u is continuous at P_0 .

The existence of ditches and ridges as well as the argument used with the comparison principle depend on the special topology of \mathbb{R}^2 . The comparison argument is similar in spirit to one used by Finn for minimal surfaces [6], and later by Finn-Giusti for constant mean curvature surfaces [8] to obtain interior gradient bounds in two dimensions.

Because $\partial\Omega$ is Lipschitz near P_0 there is a Lipschitz function $y = \phi(x)$ and a neighborhood V of the origin so that after a translation and rotation

$$(4) \quad \begin{aligned} P_0 &= (0,0) \\ V \cap \Omega &= V \cap \{P = (x,y) \text{ s.t. } y < \phi(x)\}. \end{aligned}$$

If Ω satisfies (i), (ii), or (iii) of the theorem stated in the introduction, we may also suppose that

$$(5) \quad \lim_{x \rightarrow 0^-} \phi'(x) > 0 > \lim_{x \rightarrow 0^+} \phi'(x)$$

and

$$(6) \quad \begin{aligned} \sup |\phi'(x)| &= L_1 < \tan \gamma \\ (x, \phi'(x)) &\in V. \end{aligned}$$

(The limits and sup are taken over those values of x for which $\phi'(x)$ exists).

We prove:

Theorem 1. If (4)-(6) are satisfied and if u is the variational solution to the capillary problem with contact angle γ in Ω , then u extends continuously to P_0 .

The first step is:

Lemma 1. If (4)-(6) are satisfied, there is a neighborhood $U \subset V$ of P_0 so that u is uniformly bounded in $U \cap \Omega$.

Proof. We introduce the concept of an internal sphere condition with contact angle (due to Finn-Gerhardt [7]). Domains satisfying this condition have uniformly bounded solutions u . We then show that if Ω satisfies (4)-(6) of Section 3, it satisfies the appropriate internal sphere condition with contact angle γ , near P_0 .

Definition. Let $0 < \gamma_0 < \pi/2$ and $P \in \Omega$. P satisfies an internal sphere condition with contact angle γ_0 and radius ρ ($P \in \text{i.s.c.}_{\rho, \gamma_0}$) if there is a ball B_ρ of radius ρ , $P \in \bar{B}_\rho$, such that any lower hemisphere lying above \bar{B}_ρ makes contact angle $< \gamma_0$ at all points of contact with $\partial\Omega \times \mathbb{R}$ for which the normal to $\partial\Omega$ is defined.

Definition. A neighborhood $U \cap \Omega$ satisfies a uniform internal sphere condition with contact angle γ_0 and radius δ ($U \cap \Omega \subset \text{I.S.C.}_{\delta, \gamma_0}$) if $P \in U \cap \Omega \Rightarrow \exists \rho > \delta$ s.t. $P \in \text{i.s.c.}_{\rho, \gamma_0}$.

This concept is illustrated in Figure 4. We have:

Step 1([7]). If $U \cap \Omega \subset \text{I.S.C.}_{\delta, \gamma_0}$, then any solution u to the variational capillary problem (3) with $|v| < \cos \gamma_0$ is uniformly bounded in $U \cap \Omega$.

Proof. (See [7]). This is a straightforward application of the comparison principle. The comparison surfaces are the graph of the solution, sufficiently high lower hemispheres (for the upper bound) and sufficiently low upper hemispheres (for the lower bound).




Figure 1:

The internal sphere condition with contact angle.

A corner with interior angle θ satisfies a uniform internal sphere condition for any

$$\gamma \geq \frac{\pi - \theta}{2},$$

i.e.

$$\theta \geq \pi - 2\gamma.$$

Step 2. We show that if Ω satisfies (4)-(6) there is a neighborhood U of P_0 and a $\delta > 0$ so that $U \cap \Omega$ I.S.C. $_{\delta, \gamma}$. Let $\epsilon, \rho > 0$. Take

$$U = B_\rho(P_1), \quad P_1 = (0, -\rho(1-\epsilon)).$$

For small enough ϵ, ρ , we will have $U \cap \Omega \subset \text{I.S.C.}_{\delta, \gamma}$:

For the lower hemisphere above U , $z = v(P)$, we have

$$Tv(P) = \frac{1}{\rho} (P - P_1).$$

The exterior normal n to $\partial\Omega$ is

$$n = \frac{1}{\sqrt{1 + (\phi')^2}} (-\phi', 1)$$

so that for $P = (x, y) \in B_\rho(P_1) \cap \partial\Omega$,

$$(7) \quad Tv \cdot n = \frac{1}{\rho \sqrt{1 + (\phi')^2}} (-x\phi' + y + \rho(1-\epsilon))$$

If $\phi'(x)$ exists then

$$y = \int_0^x \phi'(t) dt = \int_0^x (\phi'(t) - \phi'(x)) dt + x\phi'(x).$$

But (5) implies that the right and left-hand limits of ϕ' exist at $x = 0$. Thus, for any $\epsilon_1 > 0 \exists \rho_0 > 0$ s.t.

$$(8) \quad \rho < \rho_0 \Rightarrow y - x\phi'(x) > -\epsilon_1 \rho.$$

Combining (7) and (8) implies that if $\rho < \rho_0$,

$$(9) \quad Tv \cdot n > \frac{1 - \epsilon - \epsilon_1}{\sqrt{1 + (\phi')^2}}.$$

From (6) we have

$$(10) \quad \frac{1}{\sqrt{1 + (\phi')^2}} > \frac{1}{\sqrt{1 + L_1^2}} > \frac{1}{\sqrt{1 + \tan^2 \gamma}} = \cos \gamma,$$

so that we may choose ϵ, ϵ_1 small enough to make

$$\frac{1}{\sqrt{1 + (\phi')^2}} (1 - \epsilon - \epsilon_1) > \cos \gamma.$$

Thus. for $\rho < \rho_0(\epsilon_1)$,

$$Tv \cdot n > \cos \gamma.$$

Q.E.D.

We use the following facts often and group them as:

Lemma 2. Let U be an open set, $v \in C^2(U) \cap H^{1,1}(U)$. Then for almost all z , $\{P: v(P) = z\}$ is a regular set (a collection of simple closed curves and curves without endpoints in U) with finite total length.

If Γ is any smooth curve in U , then almost all level sets of v in U are transversal to Γ .

Proof. Almost all level sets are regular by Sard's Theorem. The co-area formula [9] implies that almost all the lengths are finite, since $v \in W^{1,1}(U)$.

Parameterize Γ by arclength s and consider the restriction of v to Γ . By Sard's theorem almost all level sets S are regular with $v'(s) \neq 0$ on S . These sets S must arise from transverse intersections between Γ and level sets of v in U .

Q.E.D.

Lemma 3. If u is bounded and (6) is satisfied (in particular, if (4)-(6) are satisfied), then given $\epsilon > 0$, $\exists \rho_1 > \rho$ s.t. for all $0 < \rho < \rho_1$,

(a) There is a ridge \bar{R}_ρ^ϵ , a connected component of the set on which $u > U - \epsilon$ satisfying $P_0 \in \bar{R}_\rho^\epsilon$ and $\bar{R}_\rho^\epsilon \cap S_\rho(P_0) \cap \Omega \neq \emptyset$.

(b) There is a ditch \bar{D}_ρ^ϵ , a connected component of the set on which $u < L + \epsilon$ satisfying $P_0 \in \bar{D}_\rho^\epsilon$ and $\bar{D}_\rho^\epsilon \cap S_\rho(P_0) \cap \Omega \neq \emptyset$.

Proof. Since u is bounded in $U \cap \Omega$, $H(S_U)$ is bounded

(1) and we may pick $M_1 < \infty$ so that

$$(11) \quad |\operatorname{div} Tu(P)| < M_1 \quad P \in U \cap \Omega.$$

Let $\epsilon > 0$ be given. To find \bar{R}_ρ^ϵ we compare S_U to the upper half of a horizontally inclined cylinder. Its axis of symmetry is parallel to the x -axis and it has radius $\frac{1}{M_1}$ (see Figure 2). Specifically, let

$$v(x, y) = \sqrt{\left(\frac{1}{M_1}\right)^2 - \left(y - \frac{1}{M_1} + \delta\right)^2} \quad \delta > 0.$$

By a useful abuse of notation we continue to call this function v and its graph S_v , even if we raise it or lower it. By construction, for all P in the domain of v and in $U \cap \Omega$,

$$(12) \quad \operatorname{div} Tv = 2H(S_v) = -M_1 < \operatorname{div} Tu.$$

Making δ small enough and picking a small enough neighborhood $U_1 \subset U$ of P_0 we can satisfy

$$(13) \quad Tv \cdot n > \cos \gamma \quad \text{on } \partial\Omega \cap U_1.$$




Figure 2:
Ridges and Ditches.

This is because $L_1 < \tan \gamma$:

$$Tv \cdot n = \frac{v_y}{\sqrt{1+v_y^2}} \frac{1}{\sqrt{1+\phi'^2}} > \frac{v_y}{\sqrt{1+v_y^2}} \frac{1}{\sqrt{1+L_1^2}} > \frac{v_y}{\sqrt{1+v_y^2}} \cos \gamma.$$

(By making δ small enough, $\frac{v_y}{\sqrt{1+v_y^2}}$ can be made arbitrarily close to 1 near $P_0 = (0,0)$).

Lift S_v so that the points P in $U_1 \cap \Omega$ above which S_v intersects S_u form a regular set with finite total length (Lemma 2), and so that

$$U - \frac{2\varepsilon}{3} < v(P_0) < U - \frac{\varepsilon}{3}.$$

Since v is continuous at P_0 there is a $\rho_1 > 0$,

$$B_{\rho_1}(P_0) \subset U_1 \text{ and}$$

$$(14) \quad U - \frac{3}{4}\varepsilon < v(P) < U - \frac{\varepsilon}{4} \quad P \in B_{\rho}(P_0)$$

For $\rho < \rho_1$, (12), (13), and the comparison principle imply that every component O in $B_{\rho} \cap \Omega$ of the set on which $u > v$ must have limit points intersecting S_{ρ} (in a set of non-zero Hausdorff measure).

Because of (14) there are points P arbitrarily near P_0 where $u(P) > v(P)$. For each $i \in \mathbb{N}$ (with $1/i < \rho$) pick P_i s.t.

$$|P_0 - P_i| < \frac{1}{i}, \quad u(P_i) > v(P_i).$$

Let O_i be the component in $B_{\rho} \cap \Omega$ of the set on which $u > v$ containing P_i . Since $\bar{O}_i \cap S_{\rho}(P_0) \neq \emptyset$, it follows that

(15)

$$|\partial O_1| > 2|\rho - \frac{1}{1}|.$$

Since $\partial(UO_1)$ has finite length in B_ρ and since at most two O_1 's can share any given boundary curve, there are only a finite number of distinct O_1 's. Hence, one of them, O_{1_1} , contains infinitely many P_1 's, so has P_0 as a limit point. From (14) it follows that O_{1_1} is part of a larger component in $B_\rho \cap \Omega$ of the set on which $u > U - \varepsilon$. This component R_ρ^ε is our ridge.

Finding the ditch D_ρ^ε uses the analogous procedure: Instead of the upper half of the cylinder placed to intersect $P_0 \times \mathbb{R}$ at a height just beneath U , one places the lower half of the cylinder so that it intersects $P_0 \times \mathbb{R}$ just above L (see Figure 1).

Q.E.D.

Remark. Lemma 3 uses the fact that $\Omega \subset \mathbb{R}^2$ crucially; the inequality (15) would not follow in general \mathbb{R}^n .

Lemma 4. Suppose (4)-(6) are satisfied. Let the following be given

$$\delta_1, \delta_2 > 0$$

$$L, U, M \text{ satisfying } -M < L < U < M.$$

Then there is a neighborhood U of $P_0 = (0,0)$ and a comparison function v defined on $U \cap \Omega$ satisfying

$$(i) \quad v \in C^2(U \cap \Omega), \quad |\operatorname{div} Tv - \kappa v| < \kappa \delta_1;$$

$$(ii) \quad \partial(U \cap \Omega) = \bigcup_{i=1}^4 \Gamma_i, \quad \text{where the } \Gamma_i \text{ are described in}$$

Figure 3, and

$$Tv \cdot n > \cos \gamma \text{ on } \Gamma_1.$$

$$v = M \text{ on } \Gamma_2$$

$$v = -M \text{ on } \Gamma_3$$

$$Tv \cdot n < \cos \gamma \text{ on } \Gamma_4.$$

$$(iii) \text{ For } P \text{ near enough } P_0, \quad |v(P) - \frac{L+U}{2}| < \delta_2.$$

Proof. (see Figure 3) S_v will be almost vertical and its level sets at height z will almost be arcs with curvature κz . (The fact that almost vertical surfaces with almost circular level sets can almost satisfy the capillary equation has been used effectively to study capillary surface behavior above corners with interior angle $\theta < \pi - 2\gamma$ [3]).

The function $z = v(x, y)$ is given implicitly (after a rotation about the z -axis) by

$$(16) \quad x = \frac{\kappa z}{2} y(y+h) + \delta(y+h)\varphi(z)$$

where h is small, $\delta < h$, and $\varphi(z)$ satisfies

$$|\varphi|_{C^2[-M, M]} < 1, \quad \frac{\partial \varphi}{\partial z} < 0, \quad \varphi = 0 \text{ in a small neighborhood of } z = \frac{L+U}{2}.$$

If $\delta = 0$, the surface S_v has a level set at height z that is part of the parabola through $(0, 0)$ and $(0, -h)$, and that has curvature almost equal to κz . For $0 < \delta < h$ the second term on the right of (16) is a small perturbation added so that v satisfies (ii) and (iii).





Figure 3:

The comparison surface S_v and its projection in \mathbb{R}^2 .



- $\Gamma_1: Tv \cdot n \geq \cos \gamma$
- $\Gamma_2: v = M$
- $\Gamma_3: v = -M$
- $\Gamma_4: Tv \cdot n \leq \cos \gamma$

Figure 2 shows the surface S_v satisfying all three conditions of Lemma 2. The key property of the differential equation (1) that allows the construction of S_v is that the mean curvature is strictly increasing with height. The key property of $\partial\Omega$ is the "half C^1 " condition (5). (It's necessary for (ii)). When it doesn't hold, for example at a re-entrant corner, u may fail to be continuous [13].

The calculations verifying (i), (ii), and (iii) of Lemma 4 are included as Appendix A. Q.E.D.

Lemma 5. (A crossing lemma in two dimensional topology)

Consider a simple closed curve Σ with interior O . Let A, B, C, D be a distinct, consecutively-ordered points on Σ . If a simple curve Γ in O has A and C as its two endpoints, then the following hold:

- (i) There is no curve in O , disjoint from Γ , connecting B to D .
- (ii) No connected open subset of O disjoint from Γ can have both B and D as limit points.

Proof. This lemma is a consequence of the Jordan Curve Theorem: Consider the simple curve obtained by following Γ from A to C , then following Σ from C to D to A . All points in a neighborhood B of B are exterior to this curve. (They can be connected to infinity). All points in a neighborhood D of D and inside O are interior to the curve (since the points outside O are exterior to it). Hence, any

path in O connecting points in $D \cap O$ to points in $B \cap O$ must cross Γ . This proves (i) and (ii). Q.E.D.

We are now in a position to prove Theorem 1. We suppose $U > L$, equations (4), (5), (6), and get a contradiction. Let

$$(17) \quad \epsilon < \frac{U-L}{6}.$$

By Lemma 3, $\exists \rho_1 > 0$ s.t. for $\rho < \rho_1$ there is a ridge R_ρ^ϵ and a ditch D_ρ^ϵ . Pick

$$(18) \quad R_\rho \in \bar{R}_\rho^\epsilon \cap S_\rho(P_0) \cap \Omega \quad D_\rho \in \bar{D}_\rho^\epsilon \cap S_\rho(P_0) \cap \Omega$$

We assume that there are arbitrarily small $\rho > 0$ for which R_ρ is oriented clockwise in Ω from D_ρ , on $S_\rho(P_0)$. This is no loss of generality: One could reflect the domain Ω and the solution u across the y -axis, preserving (4), (5), (6), and the resulting lemmas, but changing the relative orientations of D_ρ and R_ρ . Let

$$(19) \quad \delta_1 = \delta_2 = \epsilon.$$

Pick a comparison surface S_v satisfying Lemma 4. U will be the domain from this lemma, above which v is defined. From (17)-(19), and (iii) of Lemma 4, $\exists \rho_2 < \rho_1$ s.t. for $\rho < \rho_2$,

$$(20) \quad \begin{aligned} P \in \bar{R}_\rho^\epsilon \cap \Omega &\Rightarrow u(P) > U - \epsilon > \frac{U+L}{2} + 2\epsilon > v(P) + \epsilon \\ P \in \bar{D}_\rho^\epsilon \cap \Omega &\Rightarrow u(P) < L + \epsilon < \frac{U+L}{2} - 2\epsilon < v(P) - \epsilon. \end{aligned}$$

Fix a $\rho < \rho_2$ and the resulting $R_\rho^\epsilon, D_\rho^\epsilon, R_\rho, D_\rho$, with R_ρ oriented clockwise in Ω from D_ρ , on $S_\rho(P_0)$. Equations (18), (20) imply that $\bar{R}_\rho^\epsilon \cap \Omega$ is contained in a component R in $U \cap \Omega$ of the set on which $u > v + \epsilon$. Also, $\bar{D}_\rho^\epsilon \cap \Omega$ is contained in D , a component in $U \cap \Omega$ of the set on which $u < v - \epsilon$. Combining (i) of Lemma 4 with (19), (20) yields

$$(21) \quad \begin{aligned} &\text{for points in } R, \operatorname{div} T(v+\epsilon) = \operatorname{div} T v < \kappa(v+\delta_1) < \kappa u = \operatorname{div} T u \\ &\text{for points in } D, \operatorname{div} T(v-\epsilon) > \operatorname{div} T u. \end{aligned}$$

Thus, the combination of the comparison principle and all four conditions from (ii) of Lemma 4 imply

$$(22) \quad \begin{aligned} |\bar{R} \cap (\Gamma_3 \cup \Gamma_4)| &\neq 0 \\ |\bar{D} \cap (\Gamma_1 \cup \Gamma_2)| &\neq 0. \end{aligned}$$

Because of the way they were picked, (22) implies that R and D must "cross" each other (see Figure 4). But R and D cannot intersect. Making this argument rigorous is technically tedious, but proves Theorem 1.

We can assume that $\partial R, \partial D$ are regular, of finite length in $U \cap \Omega$, and that ∂R and ∂D meet $S_\rho(P_0)$ transversally. (If not, use Lemma 1 and pick a slightly larger $\epsilon < \frac{L+U}{6}$. The old R, D will be contained in a new R and D , (larger) components of the sets on which $u > v + \epsilon$, $u < v - \epsilon$, respectively).

We show that (22) cannot be satisfied.




Figure 4:

\mathcal{R} and \mathcal{D} must cross but can't.

Let R_ρ be the component of $R \cap B_\rho(P_0)$ which contains R_ρ in its closure. Let D_ρ be the component of $D \cap B_\rho(P_0)$ containing D_ρ in its closure (figure 4).

By construction $R_\rho^\epsilon \subset R_\rho$ and $D_\rho^\epsilon \subset D_\rho$ so that

$$(23) \quad P_0 \in \bar{R}_\rho, \quad P_0 \in \bar{D}_\rho.$$

There are only a finite number of other components of $R \cap B_\rho(P_0)$ having P_0 and points on $S_\rho(P_0) \cap \Omega$ as limit points, since \mathcal{R} has finite length. If any of these have limit points on $S_\rho(P_0)$ that are between R_ρ and D_ρ but nearer to D_ρ than the limit points of R_ρ , relabel: Take the one of these with the nearest (counterclockwise from D_ρ) limit points on $S_\rho(P_0)$ and call it R_ρ . Relabel one of its limit points on $S_\rho(P_0) \cap R_\rho$ as R_ρ . (Such limit points in R_ρ exist because \mathcal{R} is transversal to S_ρ). Equation (23) still holds and we proceed.

Let R_1 be the point between R_ρ and D_ρ on $S_\rho(P_0)$ nearest D_ρ and in \bar{R}_ρ . Let D_1 be the point between R_ρ and D_ρ on $S_\rho(P_0)$ nearest R_ρ and in \bar{D}_ρ (Figure 3).

We follow the curves Γ_R of \mathcal{R} through R_1 and Γ_D of \mathcal{D} through D_1 . Let Γ_R^+, Γ_D^+ be the rays initially entering $B_\rho(P_0)$. Let Γ_R^-, Γ_D^- be the rays initially leaving it.

Follow Γ_R^+ into $B_\rho(P_0)$. It cannot intersect $S_\rho(P_0)$ again. (We use (23) and Lemma 5 repeatedly here): if it intersects S_ρ

- (a) between Γ_1 and R_ρ , then $P_0 \notin \bar{R}_\rho$,
- (b) between R_ρ and R_1 or at R_1 then $R_\rho \notin \bar{R}_\rho$,
- (c) between R_1 and D_ρ , then R_1 is not the nearest point to D_ρ between R_ρ and D_ρ on $\bar{R}_\rho \cap S_\rho(P_0)$,
- (d) between D_ρ and Γ_4 , then $P_0 \notin \bar{D}_\rho$.

Because ∂R has finite length, Γ_R^+ has exactly one limit point. Because ∂R is regular this point is on $\partial \Omega \cap \overline{B_\rho(P_0)}$. It must be P_0 ; otherwise the combination of (23) and Lemma 5 (with the other three points being R_ρ , D_ρ , and P_0) would be contradicted.

The same reasoning shows that Γ_D^+ never intersects $S_\rho(P_0)$ after D_1 , and has P_0 as its limit point.

Follow Γ_R^- out of $B_\rho(P_0)$. It too must eventually have a limit point P_2 on $\partial(U \cap \Omega)$. Before it reaches R_2 it can intersect $S_\rho(P_0)$ a finite number of times between R_ρ and D_ρ . Any time it enters $B_\rho(P_0)$ through this arc it may:

(a) cross on another arc of $S_\rho(P_0)$ between Γ_1 and R_ρ or between Γ_4 and D_ρ . In either case, this would contradict the combination of (23) and Lemma 5.

(b) Not leave $B_\rho(P_0)$. Arguing as with Γ_R^+ it would follow that Γ_R^- 's limit point was P_0 . If its last intersection R_3 with S_ρ was

- (i) between R_ρ and R_1 , then R_1 could not be in \bar{R}_ρ ,
- (ii) between R_1 and D_ρ , then the arc of Γ_R^+ from R_3 to P_0 bounds a component of R

having P_0 and points on S_ρ between R_1 and D_ρ as limit points. This contradicts the paragraph following (23).

(c) Leave $S_\rho(P_0)$ through the same arc.

Thus, case (c) happens, so Γ_R intersects the arc of $S_\rho(P_0)$ between R_ρ and D_ρ an odd number of times.

Consider the simple closed curve Σ : from P_0 to R_1 (along Γ_R^+), from R_1 to R_2 (along Γ_R^-), then clockwise along $\partial(U \cap \Omega)$ back to P_0 . By construction, R_ρ and hence all of R is interior to Σ . Thus, (22) implies that

$$R_2 \in (\Gamma_3 \cup \Gamma_4) \setminus (\Gamma_2 \cap \Gamma_3).$$

Because Σ is transverse to $S_\rho(P_0)$ and intersects the arc between R_ρ and D_ρ an odd number of times, D_ρ and hence all of \mathcal{D} must be exterior to Σ . Thus, $\bar{\mathcal{D}}$ cannot intersect $\Gamma_1 \cup \Gamma_2$ and we have our contradiction to (22). Q.E.D.

§4. Related Questions

In what ways can Theorem 1 be generalized? It is clear from the proof that γ could be allowed to vary continuously along $\partial\Omega$, with $0 < \gamma(P) < \pi$. The differential equation could be generalized to

$$\operatorname{div} Tu = f(x,u), \quad f \text{ continuous in } (x,u), \quad \frac{\partial f}{\partial u} > \varepsilon_0 > 0,$$

provided it was the Euler equation for a variational problem that implies the weak boundary behavior needed for the comparison principle.

Are there situations for which $\partial\Omega$ is not regular but u still is more than just continuous?

Conjecture. If Ω is (locally) convex in a neighborhood of $P_0 \in \partial\Omega$, with Lipschitz constant $L_1 < \tan \gamma$, then u is Lipschitz continuous there.

One cannot expect much more than Lipschitz continuity in this case: One can construct a convex domain in R^2 for which P_0 is the limit (from both sides of $\partial\Omega$) of points on $\partial\Omega$ at which $\partial\Omega$ is locally a convex corner. If u is C^1 , then the contact angle condition on each segment of the corner uniquely determines the tangent plane at the corner. If $\partial\Omega$ does not have a tangent at P_0 , the limiting values of the gradients of u at these corners will be different from each side (provided $\gamma \neq \pi/2$).

Do the results of Theorem 1 generalize to R^n ? The method does not seem to.

ACKNOWLEDGMENT

The author wishes to thank Robert Finn for posing the problem studied in this paper.

Appendix A: Proof of Lemma 4

We consider the function

$$(1) \quad x = \frac{\kappa z}{2} y(y+h) + \delta(y+h)\varphi(z) = g(y,z)$$

and its graph S . We show that for a correct choice of the parameters h , δ , and the function φ , (a rotation of) S will satisfy the conditions of Lemma 4.

Require initially:

$$(a) \quad 0 < \delta < h,$$

$$(b) \quad |\varphi|_{C^2[-M,M]} < 1, \quad \varphi'(z) \leq 0$$

$$(c) \quad |z| < M, \quad -h < y < \min(\alpha|x|, h),$$

where $\alpha > 0$ is chosen large enough to guarantee that the rotation of S needed to satisfy the contact angle conditions (ii) of Lemma 4 will keep S lying above a satisfactory neighborhood $U \cap \Omega$. For example, choose

$$(d) \quad \alpha > 2 \tan \left| \frac{\pi}{2} - \gamma \right|.$$

We first find a φ and bound h so that S is a C^2 graph $z = v(x,y)$ and so that condition (iii) of Lemma 4 is satisfied. Let

$$(2) \quad \varphi(z) = \begin{cases} \mu(z_1 - z)^3 & z > z_1 = \frac{L+U}{2} + \frac{\delta}{2} \\ 0 & z_1 > z > z_0 = \frac{L+U}{2} - \frac{\delta}{2} \\ \mu(z_0 - z)^3 & z_0 > z, \end{cases}$$

where $\mu > 0$ is small, so that u satisfies (b). If $(x, y, z) \in S$ and (x, y) is near P_0 , (1) implies that $\psi(z)$ is small. It follows from (2) that if (x, y) is near enough to P_0 , z lies between the bounds of (iii).

The surface S is C^2 . To show that it is a graph $z = v(x, y)$ it suffices to show that $\frac{\partial g}{\partial z} < 0$ in the entire region (c). Since

$$(3) \quad g_z = (y+h) \left[\frac{\kappa}{2} y + \delta \psi'(z) \right]$$

and since $\psi'(z) < 0$, g_z is negative for $y < 0$. For $0 < y < \min(\alpha|x|, h)$ we are more careful. From (3) we want $\psi'(z) < -\frac{\kappa y}{2\delta}$. So for $(x, y, z) \in S$, $0 < y < \min(\alpha|x|, h)$ it suffices to find

$$(4) \quad \psi'(z) < \frac{-\kappa \alpha |x|}{2\delta}.$$

We want (4) to be satisfied for small enough h . We treat the case $x < 0$, $z > z_1$. The case $x > 0$, $z < z_0$ is analogous. If $x = 0$, then $y < 0$. From (1) and (c) we have

$$0 > x = (y+h) \left[\frac{\kappa z}{2} y + \delta \psi(z) \right] > 2h \left[\frac{\kappa \text{Max}}{2} + \delta \psi(z) \right].$$

Thus

$$x[1 - \kappa \alpha h] > 2h \delta \psi(z).$$

If:

$$(e) \quad h < h_0 < \frac{1}{\kappa \alpha},$$

then

$$(5) \quad x > 2Ch\delta\phi(z), \quad C = \frac{1}{1 - h_0 M \kappa \alpha}$$

Plugging (5) and the explicit form (2) of ϕ , into (4) gives

$$-3\mu(z_1 - z)^2 < \kappa \alpha Ch(z_1 - z)^3.$$

This inequality holds near $z_1 = z$. To make it true for all $z_1 < z < M$ the bound (e) on h may be lowered if necessary.

Thus, g_z is negative in the entire region (c) and we may write $S = S_v$. To calculate the mean curvature of S we return to the parameterization $x = g(y, z)$:

$$\begin{aligned} r = g_y &= \kappa z(y + \frac{h}{2}) + \delta\phi(z) = O(h) \\ s = g_z &= \frac{\kappa}{2}y(y + h) + \delta(y + h)\phi'(z) = O(h^2) \\ (6) \quad g_{zz} &= \delta(y + h)\phi''(z) = O(h^2) \\ g_{yy} &= \kappa z \\ g_{yz} &= \kappa(y + \frac{h}{2}) + \delta\phi'(z) = O(h). \end{aligned}$$

So

$$\begin{aligned} \operatorname{div} Tg(y, z) &= \frac{(1+s^2)g_{yy} + (1+r^2)g_{zz} - 2rs g_{yz}}{(1+r^2+s^2)^{3/2}} \\ &= \kappa z + O(h^2). \end{aligned}$$

Thus $\exists h_1$ so that

$$(f) \quad 0 < h < h_1$$

implies condition (i):

$$|\operatorname{div} Tv - \kappa v| < \kappa \delta_1.$$

It remains to verify (ii). Define Γ_2 and Γ_3 to be the sets in \mathbb{R}^2 above which S has height M and $-M$, respectively. For small enough h it is easy to calculate that the restriction of (c) that $y < h$ is redundant; the (arrowhead-shaped) region A in \mathbb{R}^2 above which S is a graph is bounded by the arcs Γ_2, Γ_3 , and the wedge $y = a|x|$.

To simplify the calculations we fix S and rotate $\partial\Omega$ about $P_0 = (0,0)$ (rather than rotating S and fixing $\partial\Omega$). We continue to use ϕ for the function describing $\partial\Omega$. At a point (x,y,z) of contact between S and $\partial\Omega \times \mathbb{R}$, the (downward) normal to S is

$$\frac{1}{\sqrt{1+r^2+s^2}} (-1, r, s).$$

The (exterior) normal to $\partial\Omega \times \mathbb{R}$ is

$$\frac{1}{\sqrt{1+\phi'^2}} (-\phi', 1, 0).$$

So their dot product is

$$(7) \quad Tv \cdot n = \frac{\phi' + r}{\sqrt{1+\phi'^2} \sqrt{1+r^2+s^2}}.$$

Since h is small and $r, s = O(h)$ the correct rotation of $\partial\Omega$ will make $\phi' / \sqrt{1+\phi'^2}$ vary near $\cos \gamma$: nearly a rotation of $\pi/2 - \gamma$ radians. The choice (d) on a insures that $\phi(x) < a|x|$ for such rotations so that S is still defined above a suitable neighborhood in Ω ; define Γ_1 and Γ_3 to be

the intersection of $\partial\Omega$ with the points in A to the left and right of $x = 0$, respectively.

From (5) of Section 3,

$$(8) \quad \lim_{x \rightarrow 0^-} \phi'(x) > \lim_{x \rightarrow 0^+} \phi'(x),$$

and by construction (see (2), (6))

$$(9) \quad \lim_{x \rightarrow 0^-} r = \frac{kh}{2} (z_1 - z_0) + \lim_{x \rightarrow 0^+} r$$

$$\lim_{x \rightarrow 0^-} s = \lim_{x \rightarrow 0^+} s.$$

It follows from (7)-(9) that for h small enough there is a rotation of $\partial\Omega$ making

$$\lim_{x \rightarrow 0^-} Tv \cdot n > \cos \gamma > \lim_{x \rightarrow 0^+} Tv \cdot n.$$

Make the parameter δ of (1) (which was free until now) small enough so that all x -values of $S \cap (\partial\Omega \times \mathbb{R})$ are near enough zero to force the conditions

$$Tv \cdot n > \cos \gamma \text{ on } \Gamma_1, \quad Tv \cdot n < \cos \gamma \text{ on } \Gamma_3.$$

This finishes the verification of (ii) and thus of the entire Lemma 4.

References

- [1] Concus, P., and Finn, R. On the behavior of a capillary surface in a wedge, Proc. Nat. Acad. Sci., 63 (1969): 292-299.
- [2] _____ On capillary free surfaces in the absence of gravity, Acta Math., 132 (1974): 177-198.
- [3] _____ On capillary free surfaces in a gravitational field, Acta Math., 132 (1974): 207-223.
- [4] _____ On the height of a capillary surface, Math. Z., 147 (1976): 93-95.
- [5] Eamer, M. Esistenza, unicità e regolarità delle superfici di equilibrio nei capillari, Ann. Univ. Ferrara Sez. VII, 18 (1973): 79-94.
- [6] Finn, R. New estimates for equations of minimal surface type, Arch. Rat. Mech. Anal., 14 (1963): 337-375.
- [7] Finn, R., and Gerhardt, C. The internal sphere condition and the capillary problem, Ann. Mat. Pura Appl., 112 (1977): 13-31.
- [8] Finn, R., and Guisti, E. On nonparametric surfaces of constant mean curvature, Ann Scuola Norm. Sup. Pisa, 4 (1977): 13-31.
- [9] Federer, H. Geometric Measure Theory, Springer-Verlag, New York (1969).
- [10] Gerhardt, C. Global regularity of solutions to the capillary problem, Ann. Scuola Norm. Sup. Pisa, 3 (1976): 157-175.

- [11] Gilbarg, D., and Trudinger, N.S. Elliptic Partial Differential Equations of Second Order, Springer-Verlag, New York (1977).
- [12] Giusti, E. Minimal Surfaces and Functions of Bounded Variation, Notes on Pure Mathematics 10, Dept. of Math., I.A.S., Australian National University, Canberra (1977).
- [13] Korevaar, N. On the behavior of a capillary surface at a re-entrant corner, Pac. J. Math., 88(2) (1980): 379-385.
- [14] Korevaar, N. Capillary Surface Behavior Determined by the Bounding Cylinder's Shape, Ph.D. dissertation, Stanford University Mathematics Department (1981).
- [15] Siegel, D. Height estimates for capillary surfaces, Pac. J. Math., 88(2) (1980): 471-515.
- [16] Simon, L. Regularity of capillary surfaces over domains with corners, Pac. J. Math., 88(2) (1980): 363-377.
- [17] Simon, L., and Spruck, J. Existence and regularity of a capillary surface with a prescribed contact angle, Arch. Rational Mech. Anal., 61 (1976): 19-34.
- [18] Spruck, J. On the existence of a capillary surface with a prescribed angle of contact, Comm. Pure Appl. Math., 28 (1975): 189-200.
- [19] Taylor, J. Boundary regularity for solutions to various capillarity and free boundary problems, Comm. Part. Diff. Eq. 2(4) (1977): 323-357.
- [20] Turkington, B. Height estimates for capillary type problems, Pac. J. Math., 88(2) (1980): 517-540.

- [21] Ural'tseva, N. N. Solution to the capillary problem,
Vestnik Leningrad Univ., 19 (1973): 54-64.
- [22] Vogel, T. J. Asymmetric unbounded liquid bridges (to
appear).
- [23] Vogel, T. I. Symmetric unbounded liquid bridges (to
appear).
- [24] Wente, H.C. The symmetry of sessile and pendant drops, Pac.
J. Math. 88(2) (1980): 387-397.

NJK/jk

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER 2439	2. GOVT ACCESSION NO. AD-4124384	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) CAPILLARY SURFACE CONTINUITY ABOVE IRREGULAR DOMAINS		5. TYPE OF REPORT & PERIOD COVERED Summary Report - no specific reporting period
		6. PERFORMING ORG. REPORT NUMBER
7. AUTHOR(s) Nicholas J. Korevaar		8. CONTRACT OR GRANT NUMBER(s) DAAG29-80-C-0041 MCS-7927062, Mod. 1
9. PERFORMING ORGANIZATION NAME AND ADDRESS Mathematics Research Center, University of 610 Walnut Street Wisconsin Madison, Wisconsin 53706		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS Work Unit Number 1 - Applied Analysis
11. CONTROLLING OFFICE NAME AND ADDRESS *see below		12. REPORT DATE October 1982
		13. NUMBER OF PAGES 35
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		15. SECURITY CLASS. (of this report) UNCLASSIFIED
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES * U.S. Army Research Office National Science Foundation P.O. Box 12211 Washington, D.C. 20550 Research Triangle Park North Carolina 27709		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) capillarity, boundary regularity, irregular, continuity, contact angle.		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) There are simple domains Ω in R^2 having re-entrant corners for which the variational solution u to the capillary problem exists, is bounded, but does not extend continuously to $\partial\Omega$. How possible is it to characterize		

20. Abstract (cont.)

domains for which u must extend continuously? This paper contains the following answer:

Theorem let $P_0 \in \partial\Omega$. Let u be the (variational) solution to the capillary problem in Ω with contact angle γ , $0 < \gamma < \pi/2$. If any of (i), (ii), or (iii) hold, then u extends continuously to P_0 :

(i) Ω is convex in a neighborhood of P_0 , $\partial\Omega$ has local Lipschitz constant $L_1 < \tan \gamma$.

(ii) $\partial\Omega$ is (locally) two C^1 curves meeting at P_0 with interior angle θ satisfying $\pi - 2\gamma < \theta < \pi$.

(iii) $\partial\Omega$ is C^1 at P_0 .

END

DATE
FILMED

3-83

DTIC